

U Kell, Douglas B.; Goodacre, Royston; Neal, Mark J.
S Institute of Biological Sciences, University of Wales, Aberystwyth, SY23
3DA, UK
O Proceedings of the ERDEC Scientific Conference on Chemical and Biological
Defense Research, Aberdeen Proving Ground, Md., Nov. 14-17, 1995 (1996),
Meeting Date 1995, 183-189. Editor(s): Berg, Dorothy A. Publisher:
National Technical Information Service, Springfield, Va.
CODEN: 64OYA8
T Conference
A English

> s specimen? and l15
264666 SPECIMEN?
16 1 SPECIMEN? AND L15

> d

16 ANSWER 1 OF 1 CA COPYRIGHT 2004 ACS on STN
N 128:50824 CA
I Application of laser microprobe inductively coupled plasma mass
spectrometry for trace metal **fingerprinting** of native gold
U Chen, Zhongxing; Doherty, Will; Gregoire, D. Conrad; Harris, Don
S Dept. of Geology, Mineral Resources Division, University of Windsor,
Windsor, ON, N9B 3P4, Can.
O Current Research - Geological Survey of Canada (1997), (1997-D), 57-62
CODEN: CRGCED; ISSN: 0704-2884
B Geological Survey of Canada
T Journal
A English

E.CNT 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

> d ab

16 ANSWER 1 OF 1 CA COPYRIGHT 2004 ACS on STN
B Major, minor, and trace elements in 60 native Au samples from across
Canada were analyzed directly by laser ablation microprobe inductively
coupled plasma-mass spectrometry (LAM-ICP-MS). Diams. of laser ablation
craters ranged from 100 to 150 μ m defining the smallest sample size
determinable. The Au signal was used as an internal standard to correct for
differences in ablation yields, matrix effects and signal **drift**
during anal. Anal. results obtained over a period of seven months for the
synthetic Au alloy standard 501 demonstrated the long-term reproducibility of
the method. The elemental signatures obtained from individual
specimens have proven to be unique and reproducible, yielding data
that can be used for matching "unknowns" with their source location.

> d l4 4

4 ANSWER 4 OF 4 CA COPYRIGHT 2004 ACS on STN
N 124:81135 CA
I Correction of Mass Spectral **Drift** Using Artificial Neural
Networks
U Goodacre, Royston; Kell, Douglas B.
S Institute of Biological Sciences, University of Wales, Aberystwyth/Dyfed,
SY23 3DA, UK
O Analytical Chemistry (1996), 68(2), 271-80
CODEN: ANCHAM; ISSN: 0003-2700
B American Chemical Society
T Journal
A English

> log y
COST IN U.S. DOLLARS

SINCE FILE TOTAL
ENTRY SESSION

ULL ESTIMATED COST	122.93	124.28
ISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
A SUBSCRIBER PRICE	-7.92	-7.92

TN INTERNATIONAL LOGOFF AT 15:44:51 ON 15 NOV 2004

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:sssptaul87mxw

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 JUL 12 BEILSTEIN enhanced with new display and select options,
resulting in a closer connection to BABS
NEWS 4 AUG 02 IFIPAT/IFIUDB/IFICDB reloaded with new search and display
fields
NEWS 5 AUG 02 Capius and CA patent records enhanced with European and Japan
Patent Office Classifications
NEWS 6 AUG 02 The Analysis Edition of STN Express with Discover!
(Version 7.01 for Windows) now available
NEWS 7 AUG 27 BIOCOMMERCE: Changes and enhancements to content coverage
NEWS 8 AUG 27 BIOTECHABS/BIOTECHDS: Two new display fields added for legal
status data from INPADOC
NEWS 9 SEP 01 INPADOC: New family current-awareness alert (SDI) available
NEWS 10 SEP 01 New pricing for the Save Answers for SciFinder Wizard within
STN Express with Discover!
NEWS 11 SEP 01 New display format, HITSTR, available in WPIDS/WPINDEX/WPIX
NEWS 12 SEP 27 STANDARDS will no longer be available on STN
NEWS 13 SEP 27 SWETSCAN will no longer be available on STN
NEWS 14 OCT 28 KOREAPAT now available on STN

NEWS EXPRESS OCTOBER 29 CURRENT WINDOWS VERSION IS V7.01A, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 11 AUGUST 2004
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 14:52:08 ON 15 NOV 2004

=> file medline

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 14:52:40 ON 15 NOV 2004

FILE LAST UPDATED: 13 NOV 2004 (20041113/UP). FILE COVERS 1950 TO DATE.

On February 29, 2004, the 2004 MeSH terms were loaded. See HELP RLOAD

for details.

OLDMEDLINE now back to 1950.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the MeSH 2004 vocabulary. See <http://www.nlm.nih.gov/mesh/> and http://www.nlm.nih.gov/pubs/techbull/nd03/nd03_mesh.html for a description of changes.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> e goodacre/au

E1	1	GOODA K/AU
E2	2	GOODA S A/AU
E3	0 -->	GOODACRE/AU
E4	22	GOODACRE A/AU
E5	6	GOODACRE A M/AU
E6	1	GOODACRE ANGELA/AU
E7	10	GOODACRE B/AU
E8	2	GOODACRE B B/AU
E9	1	GOODACRE B C/AU
E10	20	GOODACRE B W/AU
E11	1	GOODACRE BRIAN/AU
E12	4	GOODACRE BRIAN W/AU

=> e goodacre royston/au

E1	4	GOODACRE R H/AU
E2	17	GOODACRE R L/AU
E3	22 -->	GOODACRE ROYSTON/AU
E4	24	GOODACRE S/AU
E5	1	GOODACRE S C/AU
E6	2	GOODACRE S H/AU
E7	2	GOODACRE S L/AU
E8	11	GOODACRE S W/AU
E9	1	GOODACRE SARA L/AU
E10	7	GOODACRE STEVE/AU
E11	1	GOODACRE STEVE W/AU
E12	14	GOODACRE T/AU

=> s e3

L1 22 "GOODACRE ROYSTON"/AU

=> d ti 1-22

L1	ANSWER 1 OF 22	MEDLINE on STN
TI	Differentiation of Micromonospora Isolates from a Coastal Sediment in Wales on the Basis of Fourier Transform Infrared Spectroscopy, 16S rRNA Sequence Analysis, and the Amplified Fragment Length Polymorphism Technique.	
L1	ANSWER 2 OF 22	MEDLINE on STN
TI	Comparison of diffuse-reflectance absorbance and attenuated total reflectance FT-IR for the discrimination of bacteria.	
L1	ANSWER 3 OF 22	MEDLINE on STN
TI	Surface-enhanced Raman spectroscopy for bacterial discrimination utilizing a scanning electron microscope with a Raman spectroscopy interface.	
L1	ANSWER 4 OF 22	MEDLINE on STN
TI	Selective detection of proteins in mixtures using electrospray ionization mass spectrometry: influence of instrumental settings and implications for proteomics.	
L1	ANSWER 5 OF 22	MEDLINE on STN
TI	Metabolomics by numbers: acquiring and understanding global metabolite data.	
L1	ANSWER 6 OF 22	MEDLINE on STN

Whole-organism fingerprinting of the genus *Carnobacterium* using Fourier transform infrared spectroscopy (FT-IR).

ANSWER 7 OF 22 MEDLINE on STN
Ultra-violet resonance Raman spectroscopy for the rapid discrimination of urinary tract infection bacteria.

ANSWER 8 OF 22 MEDLINE on STN
Metabolic profiling: pathways in discovery.

ANSWER 9 OF 22 MEDLINE on STN
Characterization of microorganisms using UV resonance Raman spectroscopy and chemometrics.

ANSWER 10 OF 22 MEDLINE on STN
Discrimination of bacteria using surface-enhanced Raman spectroscopy.

ANSWER 11 OF 22 MEDLINE on STN
Application of high-throughput Fourier-transform infrared spectroscopy in toxicology studies: contribution to a study on the development of an animal model for idiosyncratic toxicity.

ANSWER 12 OF 22 MEDLINE on STN
Explanatory optimization of protein mass spectrometry via genetic search.

ANSWER 13 OF 22 MEDLINE on STN
Rapid quantitative assessment of the adulteration of virgin olive oils with hazelnut oils using Raman spectroscopy and chemometrics.

ANSWER 14 OF 22 MEDLINE on STN
Investigating plant-plant interference by metabolic fingerprinting.

ANSWER 15 OF 22 MEDLINE on STN
Metabolic fingerprinting of salt-stressed tomatoes.

ANSWER 16 OF 22 MEDLINE on STN
Chemometric discrimination of unfractionated plant extracts analyzed by electrospray mass spectrometry.

ANSWER 17 OF 22 MEDLINE on STN
Metabolic profiling using direct infusion electrospray ionisation mass spectrometry for the characterisation of olive oils.

ANSWER 18 OF 22 MEDLINE on STN
Fluorescent amplified fragment length polymorphism probabilistic database for identification of bacterial isolates from urinary tract infections.

ANSWER 19 OF 22 MEDLINE on STN
Monitoring of complex industrial bioprocesses for metabolite concentrations using modern spectroscopies and machine learning: application to gibberellic acid production.

ANSWER 20 OF 22 MEDLINE on STN
Sample preparation in matrix-assisted laser desorption/ionization mass spectrometry of whole bacterial cells and the detection of high mass (>20 kDa) proteins.

ANSWER 21 OF 22 MEDLINE on STN
Rapid and quantitative detection of the microbial spoilage of meat by fourier transform infrared spectroscopy and machine learning.

ANSWER 22 OF 22 MEDLINE on STN
Flow-injection electrospray ionization mass spectrometry of crude cell extracts for high-throughput bacterial identification.

> s l1 and drift
5370 DRIFT
0 L1 AND DRIFT

> file ca
OST IN U.S. DOLLARS SINCE FILE TOTAL
 ENTRY SESSION
ULL ESTIMATED COST 1.14 1.35

FILE 'CA' ENTERED AT 14:54:30 ON 15 NOV 2004
SE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
LEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

copyright of the articles to which records in this database refer is
eld by the publishers listed in the PUBLISHER (PB) field (available
or records published or updated in Chemical Abstracts after December
6, 1996), unless otherwise indicated in the original publications.
ne CA Lexicon is the copyrighted intellectual property of the
merican Chemical Society and is provided to assist you in searching
atabases on STN. Any dissemination, distribution, copying, or storing
f this information, without the prior written consent of CAS, is
trictly prohibited.

FILE COVERS 1907 - 11 Nov 2004 VOL 141 ISS 21
FILE LAST UPDATED: 11 Nov 2004 (20041111/ED)

This file contains CAS Registry Numbers for easy and accurate
substance identification.

> e goodacre r/au
1 7 GOODACRE JOHN A/AU
2 1 GOODACRE PAUL R/AU
3 19 --> GOODACRE R/AU
4 1 GOODACRE ROBERT/AU
5 1 GOODACRE ROBERT L/AU
6 86 GOODACRE ROYSTON/AU
7 1 GOODACRE S/AU
8 1 GOODACRE SARA/AU
9 1 GOODACRE SARA L/AU
10 2 GOODACRE SIMON/AU
11 3 GOODACRE SIMON C/AU
12 21 GOODACRE SIMON CHARLES/AU

> s e6
3 86 "GOODACRE ROYSTON"/AU

> s drift and l3
 34318 DRIFT
4 4 DRIFT AND L3

> d ti 1-4

4 ANSWER 1 OF 4 CA COPYRIGHT 2004 ACS on STN
I On mass spectrometer instrument standardization and interlaboratory
 calibration transfer using neural networks

4 ANSWER 2 OF 4 CA COPYRIGHT 2004 ACS on STN
I Rapid analysis of multiple determinands using pyrolysis mass spectrometry
 and supervised learning with artificial neural networks - principles and
 applications

4 ANSWER 3 OF 4 CA COPYRIGHT 2004 ACS on STN
I Analyzing a sample by using a neural network to correct for measurement
 drift

4 ANSWER 4 OF 4 CA COPYRIGHT 2004 ACS on STN
I Correction of Mass Spectral **Drift** Using Artificial Neural
 Networks

ANSWER 1 OF 4 CA COPYRIGHT 2004 ACS on STN

For pyrolysis mass spectrometry (PyMS) to be exploited in areas such as the routine identification of microorganisms, for quantifying determinants in biol. and biotechnol. systems, and in the production of useful mass spectral libraries, it is paramount that newly acquired spectra be comparable to those previously collected and held in a central reference laboratory. Artificial neural networks (ANNs) and other multivariate calibration models have been used to relate mass spectra to the biol. features of interest. However, calibration models developed on one mass spectrometer cannot be used with spectra collected on a second instrument, because of the differences between the instrumental responses of both instruments. We report here that an ANN-based **drift** correction procedure can be implemented so that newly acquired spectra can be used to challenge models constructed using mass spectra collected on different instruments. Calibration samples were run on three different PyMS machines, and ANNs set up in which the inputs were the 150 machine 'a' calibration masses and the outputs were the 150 calibration masses from the machine 'b' spectra. Such associative neural networks could thus be used as signal-processing elements to effect the transformation of data acquired on one machine to those which would have been acquired on a different instrument. Therefore, for the first time PyMS could be used to acquire spectra which could usefully be compared to those previously collected and held in a data-base, irrespectively of the mass spectrometer used. The examples reported are for the quantitative assessment of the amount of lysozyme in a binary mixture with glycogen and the rapid identification down to the species level of bacteria belonging to the genus *Eubacterium*. This approach is not limited solely to pyrolysis mass spectrometry but is generally applicable to any anal. tool which is prone to deterioration in calibration transfer, such as IR, ESR, NMR and other vibrational spectroscopies, gas and liquid chromatog., as well as other types of mass spectrometry.

ANSWER 2 OF 4 CA COPYRIGHT 2004 ACS on STN

Curie-point pyrolysis mass spectrometry is a rapid, high-resolution technique which has enjoyed historical success in the operational fingerprinting and classification of microbial and other biol. systems. More recently we have exploited the anal. power of this technique together with supervised learning, based on artificial neural networks and cognate chemometric methods, for the accurate and quant. anal. of complex biol. materials, including microorganisms, fermentor broths and agricultural products. When suitable stds. are available, supervised learning is much more powerful than in unsupervised learning for the chemometric anal. of multivariate spectroscopic data. Similar neural network methods may also be used to correct for mass spectral **drift**.

ANSWER 3 OF 4 CA COPYRIGHT 2004 ACS on STN

A method and apparatus are disclosed for analyzing a sample in which a neural network is trained to correct for measurement **drift** of a given anal. instrument (e.g., a mass spectrometer). The training is done by using first and second sets of data obtained by the instrument from samples of known compns. at initial and subsequent instants of time, respectively. The trained network is used to transform data, obtained by the instrument from a sample of unknown composition at said subsequent instant of time, to an estimate of the data which would have been obtained by the instrument from that sample at the initial instant of time. The transformed data then are analyzed to analyze the sample of unknown composition. Examples are given of the use of pyrolysis mass spectrometry for the determination of lysozyme and glycogen in mixture, ampicillin in *Escherichia coli*, *Staphylococcus aureus* in mixts. with *Escherichia coli*, and identification of human isolates of *Propionibacterium acnes*.

ANSWER 4 OF 4 CA COPYRIGHT 2004 ACS on STN

For pyrolysis mass spectrometry (PyMS) to be used for the routine identification of microorganisms, for quantifying determinands in biol. and biotechnol. systems, and in the production of useful mass spectral libraries, it is paramount that newly acquired spectra be compared to those previously collected. Neural network and other multivariate

calibration models have been used to relate mass spectra to the biol. features of interest. As commonly observed, however, mass spectral fingerprints showed a lack of long-term reproducibility due to instrumental **drift** in the mass spectrometer; when identical materials were analyzed by PyMS at dates from 4 to 20 mo apart, neural network models produced at earlier times could not be used to give accurate ests. of determinand concns. or bacterial identities. Neural networks, however, can be used to correct for pyrolysis mass spectrometer instrumental **drift** itself, so that neural network or other multivariate calibration models created using previously collected data can be used to give accurate ests. of determinand concentration or the nature of bacteria (or, indeed, other materials) from newly acquired pyrolysis mass spectra. This approach is not limited solely to pyrolysis mass spectrometry but is generally applicable to any anal. tool which is prone to instrumental **drift**, such as IR, ESR, NMR and other spectroscopies, and gas and liquid chromatog., as well as other types of mass spectrometry.

=>

=> e wilkes j/au

E1	2	WILKES IAN PETER/AU
E2	1	WILKES INGO/AU
E3	34 -->	WILKES J/AU
E4	4	WILKES J B/AU
E5	3	WILKES J D/AU
E6	13	WILKES J F/AU
E7	6	WILKES J FRED/AU
E8	25	WILKES J G/AU
E9	5	WILKES J L/AU
E10	6	WILKES J M/AU
E11	13	WILKES J O/AU
E12	1	WILKES J P/AU

=> e

E13	19	WILKES J S/AU
E14	1	WILKES JAMES C/AU
E15	1	WILKES JAMES D/AU
E16	1	WILKES JAMES E/AU
E17	4	WILKES JAMES M/AU
E18	14	WILKES JAMES O/AU
E19	1	WILKES JASON/AU
E20	5	WILKES JASON J/AU
E21	1	WILKES JASON JOHN/AU
E22	1	WILKES JEFFREY/AU
E23	1	WILKES JENNIFER J/AU
E24	3	WILKES JILLIAN/AU

=> e

E25	1	WILKES JIM/AU
E26	5	WILKES JOHN/AU
E27	53	WILKES JOHN B/AU
E28	2	WILKES JOHN BARKER/AU
E29	1	WILKES JOHN C/AU
E30	1	WILKES JOHN F/AU
E31	13	WILKES JOHN G/AU
E32	4	WILKES JOHN GEORGE/AU
E33	1	WILKES JOHN LAURENCE/AU
E34	52	WILKES JOHN S/AU
E35	1	WILKES JOHN S JR/AU
E36	1	WILKES JOHN STEPHEN/AU

=> e

E37	4	WILKES JOHN STEVEN/AU
E38	2	WILKES JOHNSTON JILLIAN/AU
E39	2	WILKES JON/AU
E40	27	WILKES JON G/AU
E41	1	WILKES JON GARDNER/AU
E42	1	WILKES JONATHAN/AU

43 6 WILKES JONATHAN M/AU
44 1 WILKES JULIE/AU
45 2 WILKES K/AU
46 6 WILKES K E/AU
47 1 WILKES K TREVOR/AU
48 1 WILKES KARL A/AU

> s e39-e41

2 "WILKES JON"/AU
27 "WILKES JON G"/AU
1 "WILKES JON GARDNER"/AU
5 30 ("WILKES JON"/AU OR "WILKES JON G"/AU OR "WILKES JON GARDNER"/AU
)

> d ti 1-30

5 ANSWER 1 OF 30 CA COPYRIGHT 2004 ACS on STN
I Predicting toxic equivalence factors from ¹³C nuclear magnetic resonance
spectra for dioxins, furans, and polychlorinated biphenyls using linear
and nonlinear pattern recognition methods

5 ANSWER 2 OF 30 CA COPYRIGHT 2004 ACS on STN
I Using Simulated 2D ¹³C NMR Nearest Neighbor Connectivity Spectral Data
Patterns to Model a Diverse Set of Estrogens

5 ANSWER 3 OF 30 CA COPYRIGHT 2004 ACS on STN
I Carbon-13 COSY NMR data for quantitative structure-activity relationships
and quantitative molecular structure-property relationships of
biomolecules

5 ANSWER 4 OF 30 CA COPYRIGHT 2004 ACS on STN
I Combining NMR spectral and structural data to form models of
polychlorinated dibenzodioxins, dibenzofurans, and biphenyls binding to
the AhR

5 ANSWER 5 OF 30 CA COPYRIGHT 2004 ACS on STN
I The use of carbon thirteen nuclear magnetic resonance spectra to predict
dioxin and furan binding affinities to the aryl hydrocarbon receptor

5 ANSWER 6 OF 30 CA COPYRIGHT 2004 ACS on STN
I Chemistry in aldol complexes of metal dications: dehydration of the
bisligand species

5 ANSWER 7 OF 30 CA COPYRIGHT 2004 ACS on STN
I Comparative structural connectivity spectra analysis (CoSCoSA) models of
steroids binding to the aromatase enzyme

5 ANSWER 8 OF 30 CA COPYRIGHT 2004 ACS on STN
I Defining and using microbial spectral databases

5 ANSWER 9 OF 30 CA COPYRIGHT 2004 ACS on STN
I Microbial identification databases

5 ANSWER 10 OF 30 CA COPYRIGHT 2004 ACS on STN
I Comparative structural connectivity spectra analysis (CoSCoSA) models of
steroid binding to the corticosteroid binding globulin

5 ANSWER 11 OF 30 CA COPYRIGHT 2004 ACS on STN
I Fragmentation Chemistry of DMSO Complexes of Metal Dications

5 ANSWER 12 OF 30 CA COPYRIGHT 2004 ACS on STN
I Fragmentation and charge transfer in gas-phase complexes of divalent metal
ions with acetonitrile

5 ANSWER 13 OF 30 CA COPYRIGHT 2004 ACS on STN
I Prediction of toxic equivalency factors for dioxins, furans and PCBs using
artificial neural networks

5 ANSWER 14 OF 30 CA COPYRIGHT 2004 ACS on STN

TI Developing ¹³C NMR quantitative spectrometric data-activity relationship
 (QSDAR) models of steroid binding to the corticosteroid binding globulin
 L5 ANSWER 15 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI ¹³C NMR Quantitative Spectrometric Data-Activity Relationship (QSDAR)
 Models of Steroids Binding the Aromatase Enzyme
 L5 ANSWER 16 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Models of Polychlorinated Dibenzodioxins, Dibenzofurans, and Biphenyls
 Binding Affinity to the Aryl Hydrocarbon Receptor Developed Using ¹³C NMR
 Data
 L5 ANSWER 17 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI ¹³C NMR and Electron Ionization Mass Spectrometric Data-Activity
 Relationship Model of Estrogen Receptor Binding
 L5 ANSWER 18 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Use of ¹³C NMR Spectrometric Data To Produce a Predictive Model of
 Estrogen Receptor Binding Activity
 L5 ANSWER 19 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Electrospray mass spectrometry for mycotoxin detection and purity analysis
 L5 ANSWER 20 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Producing ¹³C NMR, Infrared Absorption, and Electron Ionization Mass
 Spectrometric Data Models of the Monodechlorination of Chlorobenzenes,
 Chlorophenols, and Chloroanilines
 L5 ANSWER 21 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Food quality indicator device to detect volatile amines
 L5 ANSWER 22 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Sample preparation and high-resolution separation of mycotoxins possessing
 carboxyl groups
 L5 ANSWER 23 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Long-term occurrence of Trypanosoma congolense resistance to diminazene,
 isometamidium and homidium in cattle at Ghibe, Ethiopia
 L5 ANSWER 24 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Determination of underivatized fumonisin B1 and related compounds by HPLC
 L5 ANSWER 25 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Multiresidue analysis of pesticides by electrospray LC-MS and LC-MS-MS
 L5 ANSWER 26 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Diffusion based solvent removal interface for coupling liquid and
 supercritical fluid chromatography to gas chromatographic detectors
 L5 ANSWER 27 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI AC corona-discharge aerosol-neutralization device adapted to liquid
 chromatography/particle beam/mass spectrometry
 L5 ANSWER 28 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Particle size distribution is not the major factor explaining variable
 analyte transmission efficiency in liquid chromatography/particle
 beam/mass spectrometry
 L5 ANSWER 29 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Determination of fumonisins B1, B2, B3 and B4 by high-performance liquid
 chromatography with evaporative light-scattering detection
 L5 ANSWER 30 OF 30 CA COPYRIGHT 2004 ACS on STN
 TI Application of the particle beam interface to high-performance liquid
 chromatography-thermal energy analysis and electron impact mass
 spectrometry for detection of non-volatile N-nitrosamines

34318 DRIFT

L6 1 DRIFT AND L5

=> d

L6 ANSWER 1 OF 1 CA COPYRIGHT 2004 ACS on STN
AN 137:228971 CA
TI Microbial identification databases
IN **Wilkes, Jon G.**; Rafii, Fatemeh; Glover, Katherine L.; Holcomb,
Manuel; Cao, Xiaoxi; Sutherland, John B.
PA The Government of the United States of America, USA
SO U.S. Pat. Appl. Publ., 41 pp.
CODEN: USXXCO
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	US 2002138210	A1	20020926	US 2001-975530	20011010
PRAI	US 2000-239549P	P	20001010		

=> s identification and drift
273269 IDENTIFICATION
34318 DRIFT

L7 528 IDENTIFICATION AND DRIFT

=> d ti 1-10

L7 ANSWER 1 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Implications of molecular genotyping of *Helicobacter pylori* isolates from different human populations by genomic fingerprinting of enterobacterial repetitive intergenic consensus regions for strain **identification** and geographic evolution

L7 ANSWER 2 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Apparatus and method for focusing and selecting ions and charged particles at or near atmospheric pressure

L7 ANSWER 3 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Development of a Plutonium Ceramic Target for the MASHA Separator

L7 ANSWER 4 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Electrospray Ionization Ion Mobility Spectrometry of Amino Acids: Ion Mobilities and a Mass-Mobility Correlation

L7 ANSWER 5 OF 528 CA COPYRIGHT 2004 ACS on STN
TI A production at high rapidity in d + Au collisions at $\sqrt{s_{NN}} = 200$ GeV

L7 ANSWER 6 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Improved **identification** of conjugated linoleic acid isomers using silver-ion HPLC separations

L7 ANSWER 7 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Basic instrumentation for Hall A at Jefferson Lab

L7 ANSWER 8 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Temperature-modulated gas sensors: selection of modulating frequencies through noise methods

L7 ANSWER 9 OF 528 CA COPYRIGHT 2004 ACS on STN
TI Status of design and construction of the Transition Radiation Tracker (TRT) for the ATLAS experiment at the LHC

L7 ANSWER 10 OF 528 CA COPYRIGHT 2004 ACS on STN
TI A straw **drift** chamber spectrometer for studies of rare kaon decays

=> s sample drift
SAMPLE IS IGNORED AS A SCOPE FOR THIS SEARCH
L8 34318 DRIFT

=> d 17 ab

L7 ANSWER 1 OF 528 CA COPYRIGHT 2004 ACS on STN
AB Biogeog. partitioning of the genome is typical of the gastric pathogen
Helicobacter pylori. Such population-specific evolution could serve as a
model for understanding host-pathogen interaction and the impact of
genetic **drift** and recombination on insular populations. With a
total of 320 isolates from six geog. regions (Japan, India, England,
Spain, Ireland, Africa, and Peru) analyzed by enterobacterial repetitive
intergenic consensus (ERIC)-based genotyping, we examined genetic affinities
among various H. pylori populations in the world. Several strain-specific
and region-specific differences were observed by ERIC-based typing.
Polymorphic ERIC patterns indicated that the ERIC sequences are in fact
dispersed in the H. pylori chromosome at different locations separated by
various distances. Phylogenetic anal. of 61 representative isolates
revealed three distinct genetic clusters populated by isolates with shared
ERIC types independent of the cag right-junction motif type and vacA
allele status. Among the notable genetic relationships were the genotypic
similarities between Irish and Japanese and between Peruvian and Japanese
isolates. Insular genotypic characteristics of Irish isolates amid
genetic similarity to East Asian, as well as North European, strains have
been once again proved in this study. Peruvian genotypes were more
similar to those of Japanese isolates than to those of Iberian or European
isolates. Given the current debate on the origin and age of present-day
H. pylori, this is a significant finding that supports the possibility of
ancient colonization of Amerindians with East Asian strains. Genotypic
data presented here will be addnl. helpful in realizing the importance of
H. pylori geog. genomics in the development of gastro-duodenal pathol.

=> s mass spect? and 18
799626 MASS
2316030 SPECT?
260887 MASS SPECT?
(MASS(W)SPECT?)
L9 1126 MASS SPECT? AND L8

=> d ti 1-10

L9 ANSWER 1 OF 1126 CA COPYRIGHT 2004 ACS on STN
TI Resolution equations for high-field ion mobility

L9 ANSWER 2 OF 1126 CA COPYRIGHT 2004 ACS on STN
TI Nanoflow LC/IMS-MS and LC/IMS-CID/MS of protein mixtures

L9 ANSWER 3 OF 1126 CA COPYRIGHT 2004 ACS on STN
TI Development of Ion **Drift**-Chemical Ionization **Mass Spectrometry**

L9 ANSWER 4 OF 1126 CA COPYRIGHT 2004 ACS on STN
TI Construction and evaluation of polar-angle-resolved miniature
time-of-flight ion **mass spectrometer**, and its
application for electron-ion coincidence spectroscopy

L9 ANSWER 5 OF 1126 CA COPYRIGHT 2004 ACS on STN
TI Electron attachment to and anion reactions with SF5Cl: electron-swarm and
selected ion flow tube studies

L9 ANSWER 6 OF 1126 CA COPYRIGHT 2004 ACS on STN
TI Peak capacity of ion mobility **mass spectrometry**: the
utility of varying **drift** gas polarizability for the separation
of tryptic peptides

L9 ANSWER 7 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Soybean foliage residues of dicamba and 2,4-D and correlation to application rates and yield

L9 ANSWER 8 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI In situ **mass spectrometry** in a 10 Torr W chemical vapor deposition process for film thickness metrology and real-time advanced process control

L9 ANSWER 9 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI **Mass spectrometers**

L9 ANSWER 10 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Efficient direct current collision and reaction cell

=> d ti 11-20

L9 ANSWER 11 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Apparatus and method for focusing and selecting ions and charged particles at or near atmospheric pressure

L9 ANSWER 12 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Determination of thorium and uranium in Chinese foods by inductively coupled plasma **mass spectrometry**

L9 ANSWER 13 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Method and apparatus for efficient transfer of ions into a **mass spectrometer**

L9 ANSWER 14 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Time-of-flight **mass spectrometer**

L9 ANSWER 15 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Electrospray Ionization Ion Mobility Spectrometry of Amino Acids: Ion Mobilities and a Mass-Mobility Correlation

L9 ANSWER 16 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Multidynamic isotope ratio analysis using MC-ICP-MS and the causes of secular **drift** in Hf, Nd and Pb isotope ratios

L9 ANSWER 17 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI CeO₂ catalysed soot oxidation. The role of active oxygen to accelerate the oxidation conversion

L9 ANSWER 18 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Dissociation pathways of protonated water clusters

L9 ANSWER 19 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Demonstration of Proton-Transfer Reaction Time-of-Flight **Mass Spectrometry** for Real-Time Analysis of Trace Volatile Organic Compounds

L9 ANSWER 20 OF 1126 CA COPYRIGHT 2004 ACS on STN

TI Compositional analysis of isobutylene/p-methylstyrene copolymers by matrix-assisted laser desorption/ionization **mass spectrometry**

=> s fingerprint and l9

6620 FINGERPRINT

L10 2 FINGERPRINT AND L9

=> d ti 1-2

L10 ANSWER 1 OF 2 CA COPYRIGHT 2004 ACS on STN

TI Microbial identification databases

L10 ANSWER 2 OF 2 CA COPYRIGHT 2004 ACS on STN

TI Gas Sensors Arrays ('Electronic Noses'): a study about the speed/accuracy

ratio

=> d 1 ab

L10 ANSWER 1 OF 2 CA COPYRIGHT 2004 ACS on STN
AB Methods of compensating for **drift** in **fingerprint** spectra of microorganisms caused by changes in their environment are disclosed. These methods of compensating for **drift** permit identification of microorganisms from their **fingerprint** spectra regardless of the environment from which the microorganisms are obtained. Furthermore, the disclosed methods may be used to construct coherent databases of **fingerprint** spectra that may be expanded even though the standard database conditions are no longer exptl. achievable. In particular embodiments, methods of compensating for **drift** in pyrolysis **mass spectra**, constructing coherent pyrolysis **mass spectral** databases, and identifying bacteria from their pyrolysis **mass spectra** are disclosed.

=> d

L10 ANSWER 1 OF 2 CA COPYRIGHT 2004 ACS on STN
AN 137:228971 CA
TI Microbial identification databases
IN Wilkes, Jon G.; Rafii, Fatemeh; Glover, Katherine L.; Holcomb, Manuel; Cao, Xiaoxi; Sutherland, John B.
PA The Government of the United States of America, USA
SO U.S. Pat. Appl. Publ., 41 pp.
CODEN: USXXCO
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002138210	A1	20020926	US 2001-975530	20011010
PRAI	US 2000-239549P	P	20001010		

=> s specimen

L11 91655 SPECIMEN

=> d his

(FILE 'HOME' ENTERED AT 14:52:08 ON 15 NOV 2004)

FILE 'MEDLINE' ENTERED AT 14:52:40 ON 15 NOV 2004

E GOODACRE/AU

E GOODACRE ROYSTON/AU

L1 22 S E3

L2 0 S L1 AND DRIFT

FILE 'CA' ENTERED AT 14:54:30 ON 15 NOV 2004

E GOODACRE R/AU

L3 86 S E6

L4 4 S DRIFT AND L3

E WILKES J/AU

L5 30 S E39-E41

L6 1 S DRIFT AND L5

L7 528 S IDENTIFICATION AND DRIFT

L8 34318 S SAMPLE DRIFT

L9 1126 S MASS SPECT? AND L8

L10 2 S FINGERPRINT AND L9

L11 91655 S SPECIMEN

=> s 111 and 18

L12 213 L11 AND L8

> d ti 1-10

- 12 ANSWER 1 OF 213 CA COPYRIGHT 2004 ACS on STN
I Development of coincidence transmission electron microscope. III.
Incorporation with γ -type imaging energy filter
- 12 ANSWER 2 OF 213 CA COPYRIGHT 2004 ACS on STN
I Improvements in dimensional stability and lightfastness of wood by
butyrylation using microwave heating
- 12 ANSWER 3 OF 213 CA COPYRIGHT 2004 ACS on STN
I Analytical electron microscopy of InP/(In, Ga)As heterogeneous structures
- 12 ANSWER 4 OF 213 CA COPYRIGHT 2004 ACS on STN
I The use of environmental scanning electron microscopy for imaging wet and
insulating materials
- 12 ANSWER 5 OF 213 CA COPYRIGHT 2004 ACS on STN
I Investigation of the change in the **DRIFT** spectra of
light-irradiated wood with heat treatment
- 12 ANSWER 6 OF 213 CA COPYRIGHT 2004 ACS on STN
I Characteristics of carrier mobility in polyaniline
- 12 ANSWER 7 OF 213 CA COPYRIGHT 2004 ACS on STN
I Surface Modification of Colloidal Silica by Sulfonation Route for
Super-Hydrophilicity
- 12 ANSWER 8 OF 213 CA COPYRIGHT 2004 ACS on STN
I Characterization of metal-ceramic interfaces using low voltage x-ray
microanalysis in a field emission scanning electron microscope
- 12 ANSWER 9 OF 213 CA COPYRIGHT 2004 ACS on STN
I **Drift** in the hypervariable region of the hepatitis C virus
during 27 years in two patients
- 12 ANSWER 10 OF 213 CA COPYRIGHT 2004 ACS on STN
I Electron **drift** mobility in polystyrene doped with
bispyrazolopyridine derivatives

> d ti 11-20

- 12 ANSWER 11 OF 213 CA COPYRIGHT 2004 ACS on STN
I A new method for the determination of the wave aberration function for
high resolution TEM 1. Measurement of the symmetric aberrations
- 12 ANSWER 12 OF 213 CA COPYRIGHT 2004 ACS on STN
I A **specimen-drift**-free EDX mapping system in a STEM for
observing two-dimensional profiles of low dose elements in fine
semiconductor devices
- 12 ANSWER 13 OF 213 CA COPYRIGHT 2004 ACS on STN
I Automated Data Collection with a Tecnai 12 Electron Microscope:
Applications for Molecular Imaging by Cryomicroscopy
- 12 ANSWER 14 OF 213 CA COPYRIGHT 2004 ACS on STN
I Automated analysis of submicron particles by CCSEM / EDXS - where are the
limits?
- 12 ANSWER 15 OF 213 CA COPYRIGHT 2004 ACS on STN
I Automated analysis of submicron particles by computer-controlled scanning
electron microscopy
- 12 ANSWER 16 OF 213 CA COPYRIGHT 2004 ACS on STN
I Scanning probe microscope, method for measuring band structure of
substance by using the microscope, and microscopic spectroscopy
- 12 ANSWER 17 OF 213 CA COPYRIGHT 2004 ACS on STN

```

I Step bunching, step wandering and faceting. Self-organization at Si
surfaces

12 ANSWER 18 OF 213 CA COPYRIGHT 2004 ACS on STN
I Analysis of nanoscale multilayers by EDXS and EELS in the STEM

12 ANSWER 19 OF 213 CA COPYRIGHT 2004 ACS on STN
I The adsorption and curing behaviors of the epoxy/amidoamine system in the
presence of metal oxides

12 ANSWER 20 OF 213 CA COPYRIGHT 2004 ACS on STN
I A novel multicell silicon drift detector module for x-ray
spectroscopy and imaging applications

> d ab 12

12 ANSWER 12 OF 213 CA COPYRIGHT 2004 ACS on STN
B We developed a specimen-drift-free energy-dispersive
X-ray (EDX) mapping system in a scanning transmission electron microscope
(STEM) to improve the sensitivity and spatial resolution of EDX elemental
mapping images. The amount of specimen drift was
analyzed from two STEM images before and after specimen
drift by using the phase-correlation method, and was compensated
for with an image-shift deflector of the STEM by the displacement of the
scanning electron beam. We applied this system to observe the
two-dimensional distribution of low dose arsenic in silicon semiconductor
devices. The sensitivity of the elemental mapping was improved to several
tenths atomic % for arsenic atoms while maintaining a spatial resolution of 2
nm.

> d l12 and fingerprint?
AND' IS NOT A VALID FORMAT FOR FILE 'CA'
FINGERPRINT?' IS NOT A VALID FORMAT FOR FILE 'CA'

The following are valid formats:

BS ----- GI and AB
LL ----- BIB, AB, IND, RE
PPS ----- AI, PRAI
IB ----- AN, plus Bibliographic Data and PI table (default)
AN ----- List of CA abstract numbers without answer numbers
BIB ----- AN, plus Compressed Bibliographic Data
ALL ----- ALL, delimited (end of each field identified)
MAX ----- MAX, delimited for post-processing
AM ----- AN, PI and PRAI in table, plus Patent Family data
BIB ----- AN, BIB, plus Patent FAM
ND ----- Indexing data
PC ----- International Patent Classifications
AX ----- ALL, plus Patent FAM, RE
ATS ----- PI, SO
AM ----- CC, SX, TI, ST, IT
CAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;
SCAN must be entered on the same line as the DISPLAY,
e.g., D SCAN or DISPLAY SCAN)
TD ----- BIB, IPC, and NCL

ABS ----- ABS, indented with text labels
ALL ----- ALL, indented with text labels
BIB ----- BIB, indented with text labels
MAX ----- MAX, indented with text labels
STD ----- STD, indented with text labels

BIB ----- AN, plus Bibliographic Data (original)
IBIB ----- OBIB, indented with text labels

BIB ----- BIB, no citations
IBIB ----- IBIB, no citations

```


IT ----- Fields containing hit terms
 ITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)
 containing hit terms
 ITRN ----- HIT RN and its text modification
 ITSTR ----- HIT RN, its text modification, its CA index name, and
 its structure diagram
 ITSEQ ----- HIT RN, its text modification, its CA index name, its
 structure diagram, plus NTE and SEQ fields
 HITSTR ----- First HIT RN, its text modification, its CA index name, and
 its structure diagram
 HITSEQ ----- First HIT RN, its text modification, its CA index name, its
 structure diagram, plus NTE and SEQ fields
 WIC ----- Hit term plus 20 words on either side
 CC ----- Number of occurrence of hit term and field in which it occurs

to display a particular field or fields, enter the display field
 codes. For a list of the display field codes, enter HELP DFIELDS at
 the arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST;
 I,IND; TI,SO. You may specify the format fields in any order and the
 information will be displayed in the same order as the format
 specification.

all of the formats (except for SAM, SCAN, HIT, HITIND, ITRN, ITSTR,
 HITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC
 to view a specified Accession Number.

ENTER DISPLAY FORMAT (BIB):
 ENTER DISPLAY FORMAT (BIB):bib

12 ANSWER 1 OF 213 CA COPYRIGHT 2004 ACS on STN
 N 141:342470 CA
 I Development of coincidence transmission electron microscope. III.
 Incorporation with γ -type imaging energy filter
 J Nishinaka, Kenichi; Taka, Shinsuke; Kimura, Yoshihide; Takai, Yoshizo
 S Department of Material and Life Science, Graduate School of Engineering,
 Osaka University, Suita, Osaka, 565-0871, Japan
 O Journal of Electron Microscopy (2004), 53(3), 217-222
 CODEN: JELJA7; ISSN: 0022-0744
 B Oxford University Press
 T Journal
 A English

E.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

> s 112 and fingerprint?
 16668 FINGERPRINT?
 13 0 L12 AND FINGERPRINT?

> d his

(FILE 'HOME' ENTERED AT 14:52:08 ON 15 NOV 2004)

FILE 'MEDLINE' ENTERED AT 14:52:40 ON 15 NOV 2004

E GOODACRE/AU
 E GOODACRE ROYSTON/AU

22 S E3
 0 S L1 AND DRIFT

FILE 'CA' ENTERED AT 14:54:30 ON 15 NOV 2004

E GOODACRE R/AU

86 S E6
 4 S DRIFT AND L3

E WILKES J/AU

30 S E39-E41
 1 S DRIFT AND L5

528 S IDENTIFICATION AND DRIFT

34318 S SAMPLE DRIFT

1126 S MASS SPECT? AND L8

0 2 S FINGERPRINT AND L9
1 91655 S SPECIMEN
2 213 S L11 AND L8
3 0 S L12 AND FINGERPRINT?

s drift and fingerprint? amd l11
SSING OPERATOR AMD L11
e search profile that was entered contains terms or
sted terms that are not separated by a logical operator.

s drift and fingerprint? and l11
34318 DRIFT
16668 FINGERPRINT?
4 0 DRIFT AND FINGERPRINT? AND L11

s drift and fingerprint?
34318 DRIFT
16668 FINGERPRINT?
5 41 DRIFT AND FINGERPRINT?

d ti 1-10

5 ANSWER 1 OF 41 CA COPYRIGHT 2004 ACS on STN
Implications of molecular genotyping of *Helicobacter pylori* isolates from
different human populations by genomic **fingerprinting** of
enterobacterial repetitive intergenic consensus regions for strain
identification and geographic evolution

5 ANSWER 2 OF 41 CA COPYRIGHT 2004 ACS on STN
Silylation and surface properties of chemically grafted hydrophobic silica

5 ANSWER 3 OF 41 CA COPYRIGHT 2004 ACS on STN
Population genetic structure of *Titanotrichum oldhamii* (Gesneriaceae), a
subtropical bulbiliferous plant with mixed sexual and asexual reproduction

5 ANSWER 4 OF 41 CA COPYRIGHT 2004 ACS on STN
Molecular epidemiology and diagnosis of *Leishmania*: what have we learnt
from genome structure, dynamics and function?

5 ANSWER 5 OF 41 CA COPYRIGHT 2004 ACS on STN
Genetic diversity of selected bacterial populations in North Carolina

5 ANSWER 6 OF 41 CA COPYRIGHT 2004 ACS on STN
Microbial identification databases

5 ANSWER 7 OF 41 CA COPYRIGHT 2004 ACS on STN
Nonlinear electrophoresis and focusing of macromolecules

5 ANSWER 8 OF 41 CA COPYRIGHT 2004 ACS on STN
Interpretation of band differences to distinguish strains of *Serratia*
marcescens by pulsed-field gel electrophoresis of *Xba*I DNA digests

5 ANSWER 9 OF 41 CA COPYRIGHT 2004 ACS on STN
Gas Sensors Arrays ('Electronic Noses'): a study about the speed/accuracy
ratio

5 ANSWER 10 OF 41 CA COPYRIGHT 2004 ACS on STN
Low RAPD polymorphism in *Archangiopteris itoi*, a rare and endemic fern in
Taiwan

d ab 8

5 ANSWER 8 OF 41 CA COPYRIGHT 2004 ACS on STN
The number of band differences in DNA macrorestriction profiles required to
distinguish unrelated strains from an index strain varies in an outbreak
with the species and restriction enzyme used. In order to define this
difference for epidemiol. studies of *Serratia marcescens*, we produced DNA
fingerprints from 57 isolates of the organism using the

restriction enzyme XbaI and pulsed-field gel electrophoresis (PFGE). The isolates were selected on the basis of their epidemiol., serotype and phage-typing patterns to include 28 unrelated strains and 29 representatives from 2 distinct outbreaks. One of the outbreaks was prolonged, lasting for several years. Electrophoretic profiles consisting of 20 or more clearly resolved bands were obtained for all isolates. Twenty-six of the unrelated strains had unique profiles with over 10 band differences from all other strains, while 27 of the outbreak representatives could be assigned to the appropriate outbreak with confidence. The majority of the outbreak isolates had none or 2 band differences from the index profile, although 3 isolates differed by 5-7 bands. The 2 exceptions among the unrelated strains differed by 4 bands, and 3 phage typing reactions, and were isolated from London and Berlin 3 yr apart, while the 2 exceptions among the outbreak collection had clearly unique profiles with over 20 band differences from each other and the outbreak profiles. Cluster anal. using Dice coefficient and UPGMA gave cut-off values of 75-78% similarity overall for related isolates, while the closest similarity for unrelated strains was 70%. The results of this study together with those of the 6 previous reports of PFGE for *S. marcescens* (which used either enzymes XbaI or SpeI) confirm that this technique is of value for this species and that with XbaI at least, most epidemiol. related strains will only differ by 3-4 bands. However, on occasion up to 7 band differences can be found within an apparent outbreak, which may be suggestive of genetic **drift**.

> d ab 7

15 ANSWER 7 OF 41 CA COPYRIGHT 2004 ACS on STN
B Effects of nonlinear dependence **drift** velocity of (double-stranded) DNA vs. elec. field strength were investigated. In comparatively weak fields, the mol. **drift** velocity is proportional to the external elec. field, while in strong fields there is addnl. nonlinear component. This effect offers possibilities to manipulate the total **drift** velocity at will-the macromols. of different size can be made to move in opposite directions in pulsed field gel electrophoresis. A new approach for focusing DNA mols. based on nonlinear electrophoresis and geometric trapping in elec. fields is proposed. The focusing is carried out in an alternating nonuniform elec. field, created by using a wedge gel with hyperbolic boundaries. It is shown that the fractions separated in such wedge retain their rectilinear shape. Gel electrophoresis expts. supported the possibility of a pronounced nonlinear focusing of DNA mols. This nonlinear separation technique presents encouraging prospects for micromanipulating systems and also for preparative isolation of long DNA fragments and development of new separation methods for bacterial **fingerprinting**.

> d ti 11-20

15 ANSWER 11 OF 41 CA COPYRIGHT 2004 ACS on STN
I **Fingerprinting** of *Helicobacter pylori* strains by matrix-assisted laser desorption/ionization mass spectrometric analysis

15 ANSWER 12 OF 41 CA COPYRIGHT 2004 ACS on STN
I Variation of multilocus minisatellite DNA **fingerprints** in avian populations

15 ANSWER 13 OF 41 CA COPYRIGHT 2004 ACS on STN
I FTIR-spectroscopy in microbial and material analysis

15 ANSWER 14 OF 41 CA COPYRIGHT 2004 ACS on STN
I Development of an optimized sorbent for direct HPTLC-FTIR online coupling

15 ANSWER 15 OF 41 CA COPYRIGHT 2004 ACS on STN
I Channel plate for electrophoretic DNA sequencing

15 ANSWER 16 OF 41 CA COPYRIGHT 2004 ACS on STN
I Application of laser microprobe inductively coupled plasma mass

spectrometry for trace metal **fingerprinting** of native gold

L15 ANSWER 17 OF 41 CA COPYRIGHT 2004 ACS on STN
TI Separation of polyester oligomers by gradient high-performance liquid chromatography

L15 ANSWER 18 OF 41 CA COPYRIGHT 2004 ACS on STN
TI Mid-infrared as a new tool for detecting adulteration in fruit products

L15 ANSWER 19 OF 41 CA COPYRIGHT 2004 ACS on STN
TI Rapid analysis of multiple determinands using pyrolysis mass spectrometry and supervised learning with artificial neural networks - principles and applications

L15 ANSWER 20 OF 41 CA COPYRIGHT 2004 ACS on STN
TI FTIR spectroscopy

=> d 13,19 ab

L15 ANSWER 13 OF 41 CA COPYRIGHT 2004 ACS on STN
AB The investigation of the development and the properties of biofilms is difficult because classical microbiol. does not offer non-destructive methods other than microscopical observations. This paper discusses the use of different Fourier transform IR spectroscopy (FTIR-spectroscopy) techniques as a means to investigate microorganisms in biofilms. FTIR-spectroscopy is suitable for the identification of microorganisms and presents a new addition to taxonomic and genetic methods. The FTIR anal. of bacterial isolates provides **fingerprint** spectra, allowing the rapid characterization of microbial strains. Secondly, the FTIR-attenuated total reflection (ATR) technique can be used for the observation of biofilms forming directly on the interface of an ATR crystals such as germanium. These crystals can be coated to obtain a surface more relevant to study interfacial processes. Spectra can be acquired non-destructively, in situ and in real time. This method is suitable for fundamental biofilm research, as well as for monitoring of biofilm formation, e.g., in an ultrapure or drinking water systems. Furthermore, FTIR-ATR also allows the rapid anal. of deposits on surfaces, e.g., filtration membranes. The anal. discrimination between microorganisms, inorg. material or other foulants can be obtained. Thirdly, with the diffuse reflectance technique (**DRIFT**) it is possible to investigate reflecting surfaces like metals or very small samples. The composition of surface coatings like biomass or other surface contaminants can be detected. These different measurement techniques demonstrate that FTIR -spectroscopy is suitable for biofilm and surface anal. and can be applied in many different ways.

15 ANSWER 19 OF 41 CA COPYRIGHT 2004 ACS on STN
B Curie-point pyrolysis mass spectrometry is a rapid, high-resolution technique which has enjoyed historical success in the operational **fingerprinting** and classification of microbial and other biol. systems. More recently we have exploited the anal. power of this technique together with supervised learning, based on artificial neural networks and cognate chemometric methods, for the accurate and quant. anal. of complex biol. materials, including microorganisms, fermentor broths and agricultural products. When suitable stds. are available, supervised learning is much more powerful than in unsupervised learning for the chemometric anal. of multivariate spectroscopic data. Similar neural network methods may also be used to correct for mass spectral **drift**.

> d 19

15 ANSWER 19 OF 41 CA COPYRIGHT 2004 ACS on STN
N 127:106297 CA
I Rapid analysis of multiple determinands using pyrolysis mass spectrometry and supervised learning with artificial neural networks - principles and applications